

On some sequences of functions and their applications in physics

Pablo G A Braz e Silva^{†§} and Andrés R R Papa^{‡||}

[†] Department of Mathematics and Statistics, University of New Mexico, Albuquerque, NM 87131 USA

[‡] Instituto de Radioproteção e Dosimetria IRD/CNEN

Av. Salvador Allende s/n, Recreio, 22780-160 Rio de Janeiro, RJ, Brazil

Abstract. We rigorously state and prove properties previously used to illustrate non-uniqueness of some generalized physical quantities. This fact imposes peculiar constraints to the mathematical functions that can be used to represent such quantities.

1. Introduction

During the last decades many attempts have been made in order to explain and formalize mathematically a large set of phenomena that, at first glance, are not well described by traditional theories. Among those attempts we can find the whole theory of self-organized criticality (that now seems to be over at least in the criticality part) and some statistical theories. Probably the most popular in recent years has been the Daróczy-like entropy (Daróczy 1970), and its corresponding Statistical Mechanics formalism known as Tsallis entropy (Tsallis 1988):

$$S_q = k_B \frac{1 - \sum_{i=1}^W p_i^q}{q - 1}, \quad (1)$$

where W is the total number of configurations, p_i are the associated probabilities and k_B is some suitable constant. It is based in a special parameter q , which is adjusted to the phenomena the theory tries to represent. When $q \rightarrow 1$ the Boltzmann entropy is obtained. In Papa (1998), an inverse formalism was applied to find the limit when $q \rightarrow 1$. In this way, an infinite set of entropies (and consequently Statistics) of the same type was found. The necessity of a deeper criteria to choose a formalism to represent physical systems was pointed out. In particular, it was shown that for the special functional form of equation (1), integrating an infinite number of times the numerator and the denominator separately leads to the same result as differentiating once (*i.e.*, to apply the replica trick or, even simpler, L'Hôpital's rule). So, letting the number

[§] Partially supported by CAPES - BEX1901/99-0 - Brasília - Brasil

^{||} Also at International Centre for Theoretical Physics, Trieste, Italy

of integrations go to infinity, we achieve a limit that is independent of the parameter q , without considering the limit of q approaching some specific value. The aim of this paper is to state and prove a general Lemma providing general conditions to apply this technique.

2. The lemma

For a real continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ and a point $x_0 \in \mathbb{R}$, we denote by $I_n^f(x)$ the sequence of functions defined inductively as follows:

$$\begin{aligned} I_0^f(x) &= f(x) \\ I_{n+1}^f(x) &= \int_{x_0}^x I_n^f(t) dt. \end{aligned}$$

To keep the notation as simple as possible, we do not write explicitly the dependence of the sequences of functions above on the point x_0 . The point used as lower limit of integration should be obvious in what follows. We use the usual notation $f^{(j)}(x)$ to denote the j -th derivative of $f(x)$. With this basic notation, we can state the lemma:

Lemma 1 *Let $f : \mathbb{R} \rightarrow \mathbb{R}$, $g : \mathbb{R} \rightarrow \mathbb{R}$ be real functions, analytic in a neighborhood U of $x_0 \in \mathbb{R}$. Suppose $f^{(j)}(x_0) = g^{(j)}(x_0) = 0$, for $j = 0, 1, \dots, K$ and $g^{(K+1)}(x_0) \neq 0$. Let $V = [x_0 - R, x_0 + R] \subset U$, $R > 0$, be an interval where $f^{(j)}(x) \neq 0$ and $g^{(j)}(x) \neq 0$, $\forall x \in V$, $x \neq x_0$, for $j = 0, 1, \dots, K + 1$. Then,*

$$\lim_{n \rightarrow \infty} \frac{I_n^f(x)}{I_n^g(x)} = \frac{f^{(K+1)}(x_0)}{g^{(K+1)}(x_0)}$$

uniformly in $V \setminus \{x_0\}$, where the sequences of functions I_n are constructed using x_0 as the lower limit of integration.

Proof: Since f and g are analytic in V , we have for $x \in V$

$$\begin{aligned} f(x) &= \sum_{j=0}^{\infty} \frac{f^{(j)}(x_0)}{j!} (x - x_0)^j \\ g(x) &= \sum_{j=0}^{\infty} \frac{g^{(j)}(x_0)}{j!} (x - x_0)^j. \end{aligned}$$

$f^{(j)}(x_0) = g^{(j)}(x_0) = 0$, for $j = 0, 1, \dots, K$, implies

$$\begin{aligned} f(x) &= \sum_{j=K+1}^{\infty} \frac{f^{(j)}(x_0)}{j!} (x - x_0)^j \\ g(x) &= \sum_{j=K+1}^{\infty} \frac{g^{(j)}(x_0)}{j!} (x - x_0)^j. \end{aligned}$$

Since both series converge uniformly, we can integrate term by term to get

$$I_n^f(x) = \sum_{j=K+1}^{\infty} \frac{f^{(j)}(x_0)}{(j+n)!} (x - x_0)^{(j+n)}$$

$$I_n^g(x) = \sum_{j=K+1}^{\infty} \frac{g^{(j)}(x_0)}{(j+n)!} (x-x_0)^{(j+n)}$$

and then, for $x \neq x_0$,

$$\begin{aligned} \frac{I_n^f(x)}{I_n^g(x)} &= \frac{\sum_{j=K+1}^{\infty} \frac{f^{(j)}(x_0)}{(j+n)!} (x-x_0)^{(j+n)}}{\sum_{j=K+1}^{\infty} \frac{g^{(j)}(x_0)}{(j+n)!} (x-x_0)^{(j+n)}} \\ &= \frac{f^{(K+1)}(x_0) + \frac{1}{k+2+n} \left[f^{(K+2)}(x_0)(x-x_0) + \sum_{j=2}^{\infty} \frac{f^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j \right]}{g^{(K+1)}(x_0) + \frac{1}{k+2+n} \left[g^{(K+2)}(x_0)(x-x_0) + \sum_{j=2}^{\infty} \frac{g^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j \right]}. \end{aligned}$$

Since

$$\sum_{j=2}^{\infty} \frac{f^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j \quad \text{and} \quad \sum_{j=2}^{\infty} \frac{g^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j$$

converge uniformly in V , there exists a constant $C > 0$ such that for all $x \in V$,

$$\begin{aligned} \left| f^{(K+2)}(x_0)(x-x_0) + \sum_{j=2}^{\infty} \frac{f^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j \right| &\leq C \\ \left| g^{(K+2)}(x_0)(x-x_0) + \sum_{j=2}^{\infty} \frac{g^{(K+1+j)}(x_0)}{(K+1+j+n)\dots(K+3+n)} (x-x_0)^j \right| &\leq C \end{aligned}$$

and that implies

$$\lim_{n \rightarrow \infty} \frac{I_n^f(x)}{I_n^g(x)} = \frac{f^{(K+1)}(x_0)}{g^{(K+1)}(x_0)}, \quad \forall x \in V \setminus \{x_0\}.$$

3. Conclusion

We showed that under quite general conditions, the physically relevant limit can be reached from a special generalized functional form, without considering the limit to the value of the parameter that recover this point through traditional methods (replica trick or L'Hôpital's rule). This curious fact imposes very strong constraints to functions used to generalize known physical quantities. Extra conditions are necessary to choose one of them.

Acknowledgments

The authors wish to thank The International Centre for Theoretical Physics, Trieste, Italy, since their collaboration was made possible during a visit to the centre.

References

- Daróczy Z 1970 *Inf. Contro.* **16** 36
Papa A R R 1993 *J. Phys. A: Math. Gen.* **31**, 5271-5276
Tsallis C 1988 *J. Stat. Phys.* **52** 479